

DR-43

PROTIC IONIC LIQUIDS IN THE SYNTHESIS OF ANALOGS OF IMMUNOSTIMULANT PLERIXAFOR

N. K. Ratmanova,¹ I. A. Andreev,^{1,2} O. A. Ivanova,^{2,3} I. V. Trushkov^{1,2}

¹Laboratory of Chemical Synthesis, Dmitry Rogachev National Medical Research Center of Pediatric Hematology, Oncology and Immunology, 1 Samory Mashela St, Moscow, 117997, Russia;


²N. D. Zelinsky Institute of Organic Chemistry, Russian Academy of Sciences, 47 Leninsky pr., Moscow, 119991, Russia;

³M. V. Lomonosov Moscow State University, Department of Chemistry, 1–3 Leninskie gory, Moscow, 119991, Russia.

E-mail: n.ratmanova@gmail.com

Abstract. Plerixafor (also known as Mozobil®, AMD3100) is an immunostimulating agent approved by FDA in 2008 and widely used in the therapy of hematological malignancies¹. This compound is a polyamine consisting of two cyclam rings connected by a *p*-xylylene linker. The existing synthetic strategies towards plerixafor are discussed in a recent review.² To date, a variety of plerixafor structural analogs containing a *p*-xylylenediamine fragment are known. Some of them exhibit biological activity similar to that of the original drug.³

We have synthesized new structural analogs of plerixafor, in which both cyclam fragments are replaced by other nitrogen-containing moieties, such as cyclic or acyclic amines and amides. The synthetic procedure included a double donor–acceptor cyclopropane ring-opening with nitrogen-containing nucleophiles (azide, thiocyanate, *etc.*) in protic ionic liquid media acting as both a solvent and an acidic catalyst followed by post-modifications.



References

1. De Clercq E. Mozobil® (Plerixafor, AMD3100), 10 years after its approval by the US Food and Drug Administration / E. De Clercq // *Antiviral Chem. Chemother.* – 2019. – Vol. 27. – P. 1–8.
2. Ratmanova N. K. Methods for the synthesis of immunostimulant plerixafor / N. K. Ratmanova, I. A. Andreev, I. V. Trushkov // *Chem. Heterocycl. Comp.* – 2020. – Vol. 56. – P. 30–35.
3. The chemical diversity and structure-based evolution of non-peptide CXCR4 antagonists with diverse therapeutic potential / D. Peng, B. Cao, Y.-J. Zhou [et al.] // *Eur. J. Med. Chem.* – 2018. – Vol. 149. – P. 148–169.

This work was supported by the Russian Foundation for Basic Research (project # 20-33-70014) and the President of the Russian Federation scholarship for young scientists, working on R&D in top-priority areas vital to the modernization of the Russian economy (medical technologies area, # CII-278.2019.4).